

Table S2. Baseline characteristics of the patients according to sex

Characteristic	Sex		p-value ^{a)}
	Male (n=376)	Female (n=357)	
Age (y)	68.1±7.1	69.1±7.2	0.053
≥75 y	67 (17.8)	86 (24.1)	0.046
Higher education	146 (38.8)	28 (7.8)	<0.001
BMI (kg/m ²)	24.4±3.2	23.5±3.5	0.001
BMI groups			<0.001
Underweight	6 (1.6)	11 (3.1)	
Normal	102 (27.1)	152 (42.6)	
Overweight	111 (29.5)	84 (23.5)	
Obese	157 (41.8)	110 (30.8)	
Smoking	64 (17.0)	7 (2.0)	<0.001
Hypertension	291 (77.4)	284 (79.6)	0.535
Coronary artery disease	174 (46.3)	238 (66.7)	<0.001
LDL-c ≥ 1.4 mmol/L	339 (90.2)	315 (88.2)	0.470
Non-HDL-c ≥ 2.2 mmol/L	335 (89.1)	309 (86.6)	0.347
High-intensity statin therapy	70 (18.6)	94 (26.3)	0.016
Ezetimibe	9 (2.4)	18 (5.0)	0.088

Values are presented as mean±standard deviation or number (%).

BMI, body mass index; HDL-c, high-density lipoprotein cholesterol; LDL-c, low-density lipoprotein cholesterol.

^{a)}Comparisons were conducted using chi-square and Student's *t*-tests for categorical and continuous variables, respectively.