Table S1. Landmark sensitivity analysis for cardiovascular mortality: fully adjusted hazard ratios after sequentially excluding deaths within 6, 12, and 24 months of the MEC examination

Index	>6 months		>12 months		>24 months	
	aHR (95% CI)	p-value	aHR (95% CI)	p-value	aHR (95% CI)	p-value
PNI (per 1-unit)	0.92 (0.88–0.96)	<0.001	0.92 (0.87–0.96)	<0.001	0.93 (0.88–0.98)	0.011
GNRI (per 1-unit)	0.93 (0.89–0.97)	<0.001	0.93 (0.89–0.96)	<0.001	0.94 (0.90–0.97)	<0.001
CONUT (per 1-point)	1.32 (1.17–1.48)	<0.001	1.33 (1.16–1.51)	<0.001	1.30 (1.09–1.55)	0.004
log(SII), per 1-unit	1.45 (1.13–1.85)	0.003	1.40 (1.07–1.82)	0.013	1.24 (0.95–1.61)	0.114
WWI (per 1-unit)	1.14 (0.87–1.49)	0.349	1.17 (0.87–1.58)	0.289	1.12 (0.85–1.49)	0.421

PNI, Prognostic Nutritional Index; GNRI, Geriatric Nutritional Risk Index; CONUT, Controlling Nutritional Status; MEC, mobile examination center; SII, Systemic Immune-Inflammation Index; WWI, Weight-Adjusted Waist Index; aHR, adjusted hazard ratio; CI, confidence interval.

Fully adjusted Cox proportional hazards models (Model 3) refit after sequentially excluding deaths occurring within 6, 12, and 24 months of the MEC examination, pooled across 20 multiply imputed datasets using Rubin's rules.